

Pre-Op Orders General Surgery ERACS

RIH

Name

DOB

PHN

Allergies - Check (✓) one box:

-
- None known
-
- Unable to obtain
-
-
- See Allergy ADR Record

Weight (kg)

Bulleted orders are initiated by default, unless crossed out and initialed by the physician/prescriber. Boxed orders () require physician/prescriber check mark () to be initiated.

Diagnostics

- CBC, BUN, creatinine, Lytes 4, glucose, albumin, Group and Screen
- ALT, Alk Phos, bilirubin INR
- HbA1c Crossmatch _____ RBC units
- CXR ECG
- Other _____

Consults

- Anesthesiology
- Enterostomal Therapy Ostomy Referral _____
- Internal Medicine _____
- Other – initiated by office for Dr. _____

Bowel Preparation (patient to purchase required items from retail pharmacy)

- Clear fluids PO 1 day pre-op
- Oral Mechanical Bowel Preparation 1 day pre-op
- Fleet® Enema 120 mL PR at _____
- Carbohydrate Loading Pre-op Protocol
- Impact AR Protocol

Medication Instructions

- Hold antiplatelets 7 days pre-op.
- Hold warfarin 5 days pre-op. (Consider Bridging Therapy)
- Hold oral anticoagulants (e.g. dabigatran, rivaroxaban) 3 days pre-op. (Consider Bridging Therapy)
- Oral Antibiotic Protocol 1 day pre-op
- Hold the following medication for _____ days pre-op:

Other

- Skin prep as per RIH Pre-operative Skin Preparation Procedure

Day of Surgery

- BGM
- IV Ringers Lactate 15 mL/hr on an electronic infusion device

Antibiotics – Administer in OR as per IHA Recommended Drug Regimens for Surgical Prophylaxis 0–60 min pre-op

- cefazolin 2 g IV pre-op. If allergic give
 - clindamycin 600 mg IV ****AND****
 - gentamicin 1.5 mg/kg IV
- metronidazole 500 mg IV pre-op. Hold if cefazolin allergy.
- Other _____

Additional Orders

- heparin 5,000 units SC on call to OR or _____ hour(s) pre-op
- Intermittent pneumatic compression
- Other _____

Date (dd/mm/yyyy)	Time	Physician Signature	Printed Name or College ID#
/ /			