

## KETAMINE (LOW DOSE) INFUSION ORDERS

Kelowna General Hospital

Weight (kg)
-------------

Bulleted orders are initiated by default, unless crossed out and initiated by the physician/prescriber. Boxed orders ([]) require physician/prescriber check mark ([]) to be initiated.

1. ALLERGIES: See Allergy/ADR record

## 2. CODE STATUS/MOST

Refer to completed Medical Orders for Scope of Treatment (MOST) #829641

## 3. MONITORING

Document on the Acute Pain Management Flow Sheet (Adult) for monitoring (PPO #809182)

Baseline (Prior to starting infusion):

 Pulse (P), blood pressure (BP), respiratory rate (RR), sedation scale (SS), orientation to date-time-person, and pain score

## Ongoing:

- Assess Q1H × 4 hours: P, BP, RR, SS, pain score, then
- Assess Q4H and PRN: P, BP, RR, SS, pain score
- · Assess for dysphoric symptoms (illusions, unpleasant dreams, hallucinations) Q4H while awake
- · If dysphoric symptoms present give LORazepam as ordered below and page APS.

4.	MED	ICAT	ION	S

	Use Sigma Spectrum infusion pump protocol: ketamine (low dose) Infusion rate: Infuse at mg/hr (suggested range: 2 – 30 mg/hr)
_]	See Patient-Controlled Analgesia (PPO #809084) or Adult Epidural Infusion (PPO #809083) orders as per separate order set
J	LORazepam 0.5 to 1 mg SL/PO Q8H PRN (Max dose 3 mg in 24 hours) for dysphoric symptoms

	Date (dd/mm/yyyy)	Time	Physician Signature	Printed Name or College ID#
4				
	1		han	<u> </u>